

FORM PTO-1390 (Modified)
(REV 11-98)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

PP01641.102

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

Unknown 09/868293

INTERNATIONAL APPLICATION NO.
PCT/IB99/02065INTERNATIONAL FILING DATE
17 December 1999PRIORITY DATE CLAIMED
18 December 1998

TITLE OF INVENTION

CHLAMYDIA TRACHOMATIS ANTIGENS

APPLICANT(S) FOR DO/EO/US

Giulio RATTI

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ A copy of the International Search Report (PCT/ISA/210).
8. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
9. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
10. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
11. ☒ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
12. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).

Items 13 to 20 below concern document(s) or information included:

13. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☐ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☒ Certificate of Mailing by Express Mail
20. ☒ Other items or information:

Published Specification
Written Opinion

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR 1.53) 09/868293		INTERNATIONAL APPLICATION NO. PCT/IB99/02065		ATTORNEY'S DOCKET NUMBER PP01641.102	
---	--	---	--	---	--

21. The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) : <input type="checkbox"/> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1,000.00 <input checked="" type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$860.00 <input checked="" type="checkbox"/> International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$710.00 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$690.00 <input type="checkbox"/> International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00 <p style="text-align: center;">ENTER APPROPRIATE BASIC FEE AMOUNT =</p>				CALCULATIONS PTO USE ONLY <div style="border: 1px solid black; height: 100px; width: 100%;"></div>	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (e)).				\$1,570.00 \$130.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	24 - 20 =	4	x \$18.00	\$72.00	
Independent claims	2 - 3 =	0	x \$80.00	\$0.00	
Multiple Dependent Claims (check if applicable). <input checked="" type="checkbox"/>				\$270.00	
TOTAL OF ABOVE CALCULATIONS =				\$2,042.00	
Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28) (check if applicable). <input type="checkbox"/>				\$0.00	
SUBTOTAL =				\$2,042.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492 (f)).				\$0.00	
TOTAL NATIONAL FEE =				\$2,042.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable). <input type="checkbox"/>				\$0.00	
TOTAL FEES ENCLOSED =				\$2,042.00	
				Amount to be refunded	\$
				charged	\$

☒ A check in the amount of **\$2,042.00** to cover the above fees is enclosed.

☐ Please charge my Deposit Account No. _____ in the amount of _____ to cover the above fees.
 A duplicate copy of this sheet is enclosed.

☐ The Commissioner is hereby authorized to charge any fees which may be required, or credit any overpayment to Deposit Account No. **03-1664** A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Alisa A. Harbin
 CHIRON CORPORATION
 Intellectual Property -R338
 PO Box 8097
 Emeryville, California 94662-8097
 US

SIGNATURE
 Alisa A. Harbin
 NAME
 33,895
 REGISTRATION NUMBER
 June 2001
 DATE

13 AUG 2002


PATENT
Atty. Docket No. 1641.102

#6

"Express Mail" Mailing Label No. EL 701 980 365 US

Date of Deposit: August 13, 2002

I hereby certify that this paper (and all documents referred to herein) is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR §1.10 on the date indicated above and is addressed to BOX PCT, Assistant Commissioner of Patents, Washington, D.C. 20231-0001.


Esperanza C. Licad

8/13/2002
Date

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of

Ratti, Giulio

Serial No.: **09/868,293**

Group Art Unit: **To be assigned**

Filed : **02/22/2002**

Examiner: **To be assigned**

For : **CHLAMYDIA TRACHOMATIS ANTIGENS**

STATEMENT PURSUANT TO
37 C.F.R. §1.821(e), 1.821(f), 1.821(g) and 1.825(b) or 1.825(d)

Assistant Commissioner for Patents
Washington, D.C. 20231


Sir:

The undersigned hereby states that the substitute computer readable form (CRF) of the Sequence Listing is identical to the paper copy of the sequence listing submitted herewith and that no new matter has been added.

Additionally, the undersigned states that the substitute data is identical to that originally filed.

Respectfully submitted,

Dated: August 13, 2002

By: 
Rebecca M. Hale
Reg. No. 45,680

CHIRON CORPORATION
Intellectual Property
P.O. Box 8097
Emeryville, CA 94662-8097
(510) 923-3179
(510) 655-3542 (Fax)

I hereby certify that this paper is being deposited in the United States Postal Service as first class mail in an envelope addressed: BOX MISSING PARTS, Assistant Commissioner of Patents and Trademarks, Washington, D.C. 20231 February 22, 2002.

Gyne Riser
Gyne Riser

02.22.02
Date

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: GIULIO RATTI
International Filing No.: PCT/IB99/02065 Examiner: Unassigned
U.S Serial No.: 09/868,293 Group Art Unit: Unassigned
International Filing Date: December 17, 1999
For: CHLAMYDIA TRACHOMATIS ANTIGENS

DECLARATION UNDER 37 C.F.R. §1.821(f) & (g)

Assistant Commissioner of Patents
and Trademarks
Washington, D.C. 20231

Sir:

I hereby state that the sequence listing information recorded in computer readable form is identical to the paper copy of the sequence listing submitted herewith and that no new matter has been added.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.


Respectfully submitted,

Dated: Feb 22, 2002

By: Rebecca M. Hale
Rebecca M. Hale
Reg. No. 45,680

CHIRON CORPORATION
Intellectual Property - R440
P.O. Box 8097
Emeryville, CA 94608
(510) 923-3179
(510) 655-3542 (Fax)

I hereby certify that this paper is being deposited with the United States Postal Service as first class mail in an envelope addressed: BOX MISSING PARTS, Assistant Commissioner of Patents and Trademarks, Washington, D.C. 20231 on February 22, 2002.


Gyne Riser

02.22.02
Date

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: GIULIO RATTI
International Filing No.: PCT/IB99/02065 Examiner: Unassigned
U.S Serial No.: 09/868,293 Group Art Unit: Unassigned
International Filing Date: December 17, 1999
For: CHLAMYDIA TRACHOMATIS ANTIGENS

PRELIMINARY AMENDMENT

Assistant Commissioner of
Patents and Trademarks
Washington, D.C. 20231

Sir:

This Preliminary Amendment is filed in response to the Notice to Comply with Requirements for Patent Applications Containing Nucleotide Sequence and/or Amino Acid Sequence Disclosures mailed July 24, 2001. Please amend the application as follows:

In the Specification:

Pursuant to 37 C.F.R. §1.121(b), please replace the table on page 16, beginning on line 2, entitled Table III – N-TERMINAL SEQUENCES OF PROTEINS with the following rewritten table:

Spot #	N-terminal sequence
10	SKETFQRNK (SEQ ID. NO. 1)
12	TTESLETLVE (SEQ. ID NO. 2)
15	LAVSSGDQEV SQEDLLKE (SEQ. ID NO. 3)
18	XPAGNPAFPVIP (SEQ. ID NO. 4)
21	AKTRTLKGDG (SEQ ID NO. 5)
24	SDSSHNL LYNK (SEQ ID NO. 6)
25	VLLYSQASWDQ RSKADAL (SEQ. ID NO. 7)
26	KAVYVQD(A/Q)E(V/D)Q (SEQ. ID NO. 8)

31	KDxxTNGQR (SEQ ID NO. 9)
33	MSKGGQxD(Y/G) (SEQ. ID NO. 10)
38	XQXENGIVGL (SEQ. ID NO. 11)
46	MPAGNPAFPVIP (SEQ. ID NO. 12)

Please replace the table on page 16, beginning on line 7, entitled Table IV – IDENTIFICATION OF ANTIGENS with the following rewritten table:

spot	Map location	N-terminal AA seq	Annotation	CT-D gene	Predicted pI/MW
1	OMP2 cluster	-	OMP2	<i>omcB</i>	7.65-7.92/54.5-58.7
2	5.2-5.3/59.7	VA(D/K)NI(K/F)YNEE (SEQ ID NO. 13)	GroEL-like	<i>groEL1</i>	5.11/58.1
3	4.6-4.9/40	LPVGN (SEQ ID NO. 14)	MOMP	<i>ompA</i>	4.69/40.3
4	4.92-5.04/70.5	SEKRK(S/A)N(K/S). (SEQ ID NO. 15).	DnaK-like	<i>dnaK</i>	4.88/70.7
10	5.44-5.64/42.2	SKETFQRNK (SEQ ID NO. 1)	EF-Tu	<i>tufA</i>	5.36/43.1
12	4.80/15.8	TTESLETLVE (SEQ ID NO. 2)	Ribosomal protein L7/12	<i>r17</i>	5.09/13.5
15	5.89/48.4	LAVSSGDQEVSQEDLLKE (SEQ ID NO. 3)	stress induced protease	<i>htrA</i>	5.83/49.5
18	5.08/34.09	XPAGNPAFPVIP (SEQ ID NO. 4)	outer membrane protein	<i>ompB</i>	5.06/34.5
19	5.14-5.28/69	Not determined	Ribosomal protein S1	<i>rs1</i>	5.17/63.6
21	5.27/40.5	AKRTRLKGDG (SEQ ID NO. 5)	EF-Tu related peptide?	-	-
24	5.32/40.5	SDSSHNLLYNK (SEQ ID NO. 6)	RNAP alpha chain	<i>rpoA</i>	5.34/41.7
25	5.97/47.6	VLLYSQASWDQQRKADAL (SEQ ID NO. 7)	Aminopeptidase	<i>pepA</i>	5.74/54.0
26	5.68/48.6	KAVYVQD(A/Q)E(V/D)Q (SEQ ID NO. 8)	Not identified	-	-
31	5.43/40.4	KDxxTNGQR (SEQ ID NO. 9)	Not identified	-	-
33	6.64/25.4	MSKGGQxD(Y/G) (SEQ ID NO. 10)	Not identified	-	-
38	5.23/40.1	XQXENGIVGL (SEQ ID NO. 11)	GTP-binding protein	<i>ychF</i>	5.16/39.5
46	5.19/33.4	MPAGNPAFPVIP (SEQ ID NO. 12)	outer membrane protein	<i>ompB</i>	5.06/34.5

A marked-up version of the changes made to the specification is attached to this amendment.

Remarks

In addition to the instant Preliminary Amendment, a declaration in accordance with §1.825(f) & (g) is submitted to comply with Sequence Listing rules, §1.821-§1.825. In the declaration, the undersigned states that the attached Sequence Listing does not introduce new matter into the specification, and the above amendments to the specification insert references to the sequence listing, as required. Further the undersigned states that the matter of the computer-

readable format is the same as the paper sequence listing. Also accompanying this amendment is a diskette with the computer readable format.

TABLE III – N-TERMINAL SEQUENCES OF PROTEINS

Spot #	N-terminal sequence
10	SKETFQRNK (SEQ ID NO. 1)
12	TTESLETLVE (SEQ. ID NO. 2)
15	LAVSSGDQEVSQEDLLKE (SEQ. ID NO. 3)
18	XPAGNPAFPVIP (SEQ. ID NO. 4)
21	AKTRTLKGDG (SEQ ID NO. 5)
24	SDSSHNLLYNK (SEQ ID NO. 6)
25	VLLYSQASWDQRSKADAL (SEQ. ID NO. 7)
26	KAVYVQD(A/Q)E(V/D)Q (SEQ. ID NO. 8)
31	KDxxTNGQR (SEQ ID NO. 9)
33	MSKGGQxD(Y/G) (SEQ. ID NO. 10)
38	XQXENGIVGL (SEQ. ID NO. 11)
46	MPAGNPAFPVIP (SEQ. ID NO. 12)

TABLE IV – IDENTIFICATION OF ANTIGENS

“CT-D gene” refers to the gene name from reference 2 and gives the names of genes likely to encode homologue proteins in *C. trachomatis* D. Theoretical pI/MW values in the last column, to be compared to the experimental values, were calculated the from CT-D gene sequences.

spot	Map location	N-terminal AA seq	Annotation	CT-D gene	Predicted pI/MW
1	OMP2 cluster	-	OMP2	<i>omcB</i>	7.65-7.92/54.5-58.7
2	5.2-5.3/59.7	VA(D/K)NI(K/F)YNEE (SEQ ID NO. 13)	GroEL-like	<i>groEL1</i>	5.11/58.1
3	4.6-4.9/40	LPVGN (SEQ ID NO. 14)	MOMP	<i>ompA</i>	4.69/40.3
4	4.92-5.04/70.5	SEK RK(S/A)N(K/S). (SEQ ID NO. 15)	DnaK-like	<i>dnaK</i>	4.88/70.7
10	5.44-5.64/42.2	SKETFQRNK (SEQ ID NO. 1)	EF-Tu	<i>tufA</i>	5.36/43.1
12	4.80/15.8	TTESLETLVE (SEQ ID NO. 2)	Ribosomal protein L7/12	<i>r17</i>	5.09/13.5
15	5.89/48.4	LAVSSGDQEVSQEDLLKE (SEQ ID NO. 3)	stress induced protease	<i>htrA</i>	5.83/49.5
18	5.08/34.09	XPAGNPAFPVIP (SEQ ID NO. 4)	outer membrane protein	<i>ompB</i>	5.06/34.5
19	5.14-5.28/69	Not determined	Ribosomal protein S1	<i>rs1</i>	5.17/63.6
21	5.27/40.5	AKTRTLKGDG (SEQ ID NO. 5)	EF-Tu related peptide?	-	-
24	5.32/40.5	SDSSHNLLYNK (SEQ ID NO. 6)	RNAP alpha chain	<i>rpoA</i>	5.34/41.7
25	5.97/47.6	VLLYSQASWDQRSKADAL (SEQ ID NO. 7)	Aminopeptidase	<i>pepA</i>	5.74/54.0
26	5.68/48.6	KAVYVQD(A/Q)E(V/D)Q (SEQ ID NO. 8)	Not identified	-	-
31	5.43/40.4	KDxxTNGQR (SEQ ID NO. 9)	Not identified	-	-
33	6.64/25.4	MSKGGQxD(Y/G) (SEQ ID NO. 10)	Not identified	-	-
38	5.23/40.1	XQXENGIVGL (SEQ ID NO. 11)	GTP-binding protein	<i>ychF</i>	5.16/39.5
46	5.19/33.4	MPAGNPAFPVIP (SEQ ID NO. 12)	outer membrane protein	<i>ompB</i>	5.06/34.5

3/PRTS

CHLAMYDIA ANTIGENS

This invention relates to antigenic proteins from *Chlamydia trachomatis*. In particular, it relates to antigens which are recognised by antibodies from chronically infected or convalescent patient sera.

5 BACKGROUND

The *Chlamydia* are obligate intracellular parasites of eukaryotic cells which are responsible for endemic sexually transmitted infections, trachoma, infectious pneumonitis, and various other disease syndromes. They occupy an exclusive eubacterial phylogenic branch, having no close relationship to any other known organisms – they are classified in their own order
10 (*Chlamydiales*) which contains a single family (*Chlamydiaceae*) which in turn contains a single genus (*Chlamydia*). Four chlamydial species are currently known – *C.trachomatis*, *C.pneumoniae*, *C.pecorum* and *C.psittaci* [eg. see reference 1]. A genome sequence of *C.trachomatis* (serovar D) has recently been published [2].

The human serovariants (“serovars”) of *C.trachomatis* are divided into two biovariants
15 (“biovars”). Serovars A-K elicit epithelial infections primarily in the ocular tissue (A-C) or urogenital tract (D-K). Serovars L1, L2 and L3 are the agents of invasive lymphogranuloma venereum (LGV).

Although chlamydial infection itself causes disease, it is thought that, in some patients, the severity of symptoms is due, in fact, to an aberrant host immune response. Failure to clear the
20 infection results in persistent immune stimulation and, rather than helping the host, this results in chronic infection with severe consequences, including sterility and blindness [3].

In addition, the protection conferred by natural chlamydial infection, is usually incomplete, transient, and strain-specific.

Due to the serious nature of the disease, there is a desire to provide suitable vaccines. These
25 may be useful (a) for immunisation against chlamydial infection or against chlamydia-induced disease (prophylactic vaccination) or (b) for the eradication of an established chronic chlamydial infection (therapeutic vaccination). Being an intracellular parasite, however, the bacterium can generally evade antibody-mediated immune responses.

Various antigenic proteins have been described for *C.trachomatis*, and the cell surface in
30 particular has been the target of detailed research [eg. 1,4]. These include, for instance, pgp3 [5,6,7], MOMP [8], Hsp60 (GroEL) [9] and Hsp70 (DnaK-like) [10]. Not all of these have

proved to be effective vaccines. however, and it is an object of the invention to identify chlamydial antigens which elicit an immune response during natural infection, in order to provide antigens and immunogens suitable for use in vaccine development.

DESCRIPTION OF THE INVENTION

5 The invention is based on the identification of proteins encoded by *Chlamydia trachomatis* which are immunogenic in man as a consequence of infection.

The invention provides a *C.trachomatis* protein having the MW and pI characteristics of protein 5, 6, 7, 8, 9, 11, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, or 55, as set
10 out in Table II on page 15.

These include proteins having, in the L2 strain of *C.trachomatis*, an N-terminal amino acid sequence disclosed in Table III on page 16.

The invention also provides proteins having sequence identity to these *C.trachomatis* proteins. Depending on the particular protein, the degree of identity is preferably greater than 50% (eg.
15 65%, 80%, 90%, 95%, 98%, 99% or more). These homologous proteins include mutants, allelic variants, serovariants, and biovariants. Identity between the proteins is preferably determined by the Smith-Waterman homology search algorithm as implemented in the MPSRCH program (Oxford Molecular). using an affine gap search with parameters *gap open penalty=12* and *gap extension penalty=1*. Typically, 50% identity or more between two
20 proteins is considered to be an indication of functional equivalence.

The invention further provides proteins comprising fragments of the *C.trachomatis* proteins of the invention. The fragments should comprise at least *n* consecutive amino acids from the proteins and, depending on the particular protein, *n* is 7 or more (eg. 8, 10, 12, 14, 16, 18, 20, 50, 100 or more). Preferably the fragments comprise an epitope from the protein.

25 The proteins of the invention can, of course, be prepared by various means (eg. recombinant expression, purification from cell culture, chemical synthesis *etc.*) and in various forms (eg. native, fusions *etc.*). They are preferably prepared in substantially isolated or purified form (*ie.* substantially free from other *C.trachomatis* or host cell proteins)

According to a further aspect, the invention provides antibodies which bind to these proteins.

30 These may be polyclonal or monoclonal and may be produced by any suitable means.

According to a further aspect, the invention provides nucleic acid encoding the proteins and protein fragments of the invention. Nucleic acid having sequence identity to this nucleic acid is also provided. Depending on the particular nucleic acid, the degree of identity is preferably greater than 50% (eg. 65%, 80%, 90%, 95%, 98%, 99% or more).

- 5 Furthermore, the invention provides nucleic acid which can hybridise to this nucleic acid, preferably under "high stringency" conditions (eg. 65°C in a 0.1xSSC, 0.5% SDS solution).

Fragments of this nucleic acid are also provided. The fragments should comprise at least n consecutive nucleotides from the sequences and, depending on the particular sequence, n is 10 or more (eg. 12, 14, 15, 18, 20, 25, 30, 35, 40 or more).

- 10 It should also be appreciated that the invention provides nucleic acid comprising sequences complementary to those described above (eg. for antisense or probing purposes).

Nucleic acid according to the invention can, of course, be prepared in many ways (eg. by chemical synthesis, from genomic or cDNA libraries, from the organism itself *etc.*) and can take various forms (eg. single stranded, double stranded, vectors, probes *etc.*).

- 15 In addition, the term "nucleic acid" includes DNA and RNA, and also their analogues, such as those containing modified backbones, and also peptide nucleic acids (PNA) *etc.*

According to a further aspect, the invention provides vectors comprising nucleic acid of the invention (eg. expression vectors) and host cells transformed with such vectors.

- 20 According to a further aspect, the invention provides compositions comprising protein, antibody, and/or nucleic acid according to the invention. These compositions may be suitable as immunogenic compositions (including vaccines), for instance, or as diagnostic reagents.

- The invention also provides nucleic acid, protein, or antibody according to the invention for use as medicaments (eg. as vaccines) or as diagnostic reagents. In particular, the invention provides protein 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 25 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, or 55 (as set out in Table II on page 15) for use as a chlamydial immunogen. Whilst it is believed that some of the proteins described in Table II may be known *per se*, they have not been disclosed as being immunogenic.

- 30 The invention also provides the use of nucleic acid, protein, or antibody according to the invention in the manufacture of (i) a medicament for treating or preventing infection due to *Chlamydia*; (ii) a diagnostic reagent for detecting the presence of *Chlamydia* or of antibodies

raised against *Chlamydia*; and/or (iii) a reagent which can raise antibodies against *Chlamydia*. The *Chlamydia* may be any species or strain, but is preferably *C.trachomatis*. In preferred embodiments, the invention provides a protein of the 55 proteins of Table II for use in such manufacture.

- 5 The invention also provides a method of treating a patient, comprising administering to the patient a therapeutically effective amount of nucleic acid, protein, and/or antibody according to the invention.

According to further aspects, the invention provides various processes.

- 10 A process for producing proteins of the invention is provided, comprising the step of culturing a host cell according to the invention under conditions which induce protein expression.

A process for producing protein or nucleic acid of the invention is provided, wherein the protein or nucleic acid is synthesised in part or in whole using chemical means.

- 15 A process for detecting nucleic acid of the invention is provided, comprising the steps of: (a) contacting a nucleic probe according to the invention with a biological sample under hybridising conditions to form duplexes; and (b) detecting said duplexes.

- A process for detecting proteins of the invention is provided, comprising the steps of: (a) contacting an antibody according to the invention with a biological sample under conditions suitable for the formation of an antibody-antigen complexes; and (b) detecting said complexes. Similarly, the invention provides a process for detecting anti-chlamydial antibodies in a sample, comprising the steps of: (a) contacting a protein according to the invention with a biological sample under conditions suitable for the formation of an antibody-antigen complexes; and (b) detecting said complexes.
- 20

The invention also provides kits comprising reagents suitable for use in these processes.

- 25 A kit is provided comprising a nucleic probe according to the invention and means for detecting duplexes formed by the probe. A kit is provided comprising an antibody according to the invention and means for detecting antibody-antigen complexes formed by the antibody. A kit is provided comprising a protein according to the invention and means for detecting antibody-antigen complexes formed by the protein.

- 30 For the avoidance of doubt, the term "comprising" encompasses "including" as well as "consisting" eg. a composition "comprising" X may consist exclusively of X or may include something additional to X, such as X+Y.

DESCRIPTION OF THE DRAWINGS

Figure 1 shows the annotated reference 2D electrophoretic EB map, also indicating the positions of the immunoreactive protein spots, labelled 1-55. Groups of spots which appear to be an isoelectric series of the same protein are encircled together and classified under the same identification number.

Figure 2 shows typical immunoblots. The whole map area is shown. Major known immunogens are marked for easier comparison. For other spot identification, refer to Figure 1 and table II. Blot A is from PID patient JO51 (MIF titre 256), and has a serum dilution 1:5000. Blot B is from patient JO35 (MIF titre 64) affected by secondary sterility, and has a serum dilution 1:2500. Blot C is similar to blot B, but is from patient JO52. Blot D is from PID patient JO31 (MIF titre 256), and has a serum dilution 1:5000.

EXAMPLES

Human sera

Sera (Table I) were obtained from women who had responded to a chlamydial infection of the genital tract. The seventeen sera (A...Q) were obtained from 4 cases of lower genital tract infection and 13 laparoscopically-confirmed cases of PID (pelvic inflammatory disease), including 2 cases of secondary sterility. All sera were positive for a standard microimmunofluorescence test (MIF) with purified *C.trachomatis* L2 elementary bodies [11], and confirmed as *C.trachomatis* immune sera by an ELISA test with the plasmid-encoded pgp3 antigen [5].

A group of 10 seronegative control sera from healthy blood donors was tested by immunoblotting in the same way, and using the same dilutions as for patient sera, in order to exclude the occurrence of non-specific reactions.

Most sera were obtained from the Chlamydia collection of the Biobanque de Picardie (Amiens, France). Some PID and control sera from healthy blood donors were obtained from the Ospedale Policlinico S.Orsola (Bologna, Italy).

Preparation of protein samples

Purified chlamydial cells were obtained as described in reference 12, by growing *C.trachomatis* strain L2/343/Bu in Vero cell cultures according to standard procedures, followed by two cycles of density gradient centrifugation [13]. The average protein concentration of the purified elementary body (EB) preparation was determined using a biuret

assay. Aliquots (2mg protein/ml) were stored in water at -20°C for subsequent electrophoretic analysis. The cells used were mainly in the form of EBs – all known chlamydial antigens to date have been found in elementary bodies, rather than reticular bodies.

Separation of chlamydial proteins

5 Chlamydial proteins were separated using high resolution 2D electrophoresis, performed using the immobiline/polyacrylamide system, essentially as described in references 14 and 15.

For analytical gels, approximately $45\mu\text{g}$ total elementary body protein was used per gel. For semipreparative gels (for microsequencing), approximately 1mg protein was used. Aliquots of the EB proteins were pelleted by low-speed centrifugation and resuspended in 8M urea, 4%
10 CHAPS (3-[(3-cholamidopropyl)dimethylammonium]-1-propane sulfonate), 40mM Tris base, 65mM dithioerythritol (DTE) and trace amounts of bromophenol blue.

Isoelectric focusing was carried out on immobiline strips providing a non-linear 3 to 10 pH gradient (IPG strips, Amersham Pharmacia Biotech). Voltage was linearly increased from 300 to 3500 V during the first three hours, then stabilised at 5000 V for 22 hours (total Volts-hour product = 110kVh). After electrophoresis, IPG strips were equilibrated for 12 min against 6 M
15 urea, 30% glycerol, 2% SDS, 0.05 M Tris.HCl, pH 6.8, and 2% DTE. The second dimension was carried out in a Laemmli system on 9-16% polyacrylamide linear gradient gels (18 cm x 20 cm x 1.5 mm), at 40 mA/gel constant current, for approximately 5 hours until the dye front reached the bottom of the gel. Analytical gels were stained with ammoniacal silver nitrate
20 [16]. The protein maps were scanned with a laser photodensitometer (Molecular Dynamics) and converted into electronic files which were then analysed with the Melanie II computer software (Bio-Rad).

Figure 1 shows the annotated reference EB map which was used to identify proteins on immunoblots. MW and pI coordinates for the reference map were calibrated by co-migration
25 of the chlamydial proteins with human serum proteins acting as reference proteins. The isoelectric point values used for serum proteins were those described in reference 17.

Immunoblot analysis

Immunoblotting results are summarised in Figure 2 and Table II.

After two-dimensional electrophoresis, the gels were electroblotted onto nitrocellulose
30 membranes [18], and processed according to standard procedures, modified as described in reference 19. Briefly, before immunodetection, the membranes were stained in 0.2% (w/v)

Ponceau S in 3% (w/v) trichloroacetic acid for 3 minutes and the positions of selected anchor spots were marked on the blot to assist matching of the immunoblots with the silver stained map. Immunoreactive spots were detected by overnight incubation at room temperature with patient sera (1500-5000x dilutions), followed by incubation with rabbit anti-human IgGs conjugated with peroxidase (Cappel, 7000x dilution), and detection with a chemiluminescence based kit (Pharmacia Amersham Biotech).

Typically, six identical 2D maps were prepared in parallel for each experiment – five were blotted onto nitrocellulose and one was stained with silver nitrate for subsequent correlation with the immunoblots and computer-assisted matching to the reference map.

- 10 The spot signals on the immunoblot almost always corresponded to a spot on the silver stained gel. However, in at least two instances (spots 13 and 14 in Figure 1), immunoblot analysis detected protein spots which were not visible in the silver stained map. This shows that this technique has a superior sensitivity and should be taken into consideration as a valuable tool also for systematic proteomics studies.
- 15 To assist matching of the immunoblot with the reference map shown in Figure 1, the nitrocellulose blots were marked with a number of internal “anchor” spots using transient Ponceau Red staining. After incubation with the sera and detection of bound antibodies by chemiluminescence, the immunoblot images were matched to the reference map and spots were assigned the corresponding pI and MW coordinates (see Table II). When the position and
- 20 shape of the spot (or isoelectric series of spots) coincided with a previously-identified EB antigen, an immune response against such antigen was recorded. In all other cases the immunoblot spot was identified by the MW and pI coordinates taken at the baricentre of the stained area (or the coordinate range, in the case of complex spot patterns). It will be appreciated that the MW and pI values are determined electrophoretically, and may have a
- 25 potential average error of +/-10%. The higher MW measurements will tend to be less accurate.

While control blots were totally blank, patient blots showed individually different patterns comprising a number of spots, which varied from 2 to 28, with an average of around 15 (see Table II). The number of immunoreactive spots had did not correlate with the serum MIF titres (see Tables I and II), so blot patterns appear to reflect a real individual variation in humoral

30 responses, and not just the difference of antibody titres. This was also confirmed by comparing the results of each serum at various dilutions.

Typical immunoblot results are shown in Figure 2. The only constant feature for all examined sera was the presence of antibodies against a complex cluster of spots previously identified as due to the cysteine-rich outer membrane protein OMP2 [12]. This cluster is shown as spot 1 in Figure 1 – all the spots labelled “1” were scored as a single antigen, but a number of accessory spots with lower MW and pI values which usually appear associated to OMP2 reactivity were separately scored, as their relationship to the OMP2 polypeptide is still unclear. Because the OMP2 protein is chlamydia-specific, and does not seem to undergo any relevant antigenic variation, it can be considered probably the best marker of chlamydial infection in this study.

The next-most frequent spots which were observed correspond to the following:

- Spot 2 – the GroEL-like (hsp60) protein (15/17 patients)
- Spot 3 – the major outer membrane protein MOMP (13/17 patients)
- Spot 4 – the DnaK-like (hsp70) protein (11/17 patients).

Reactivity with these known immunogens can be considered as an internal control which demonstrates the quality of the human sera used in this study. The lack of pgp3 reactivity on the blots, however, is significant because all the sera had been found positive in an ELISA confirmatory assay with a purified soluble form of pgp3. This suggests that antibody response to pgp3 in human infections occurs mainly against epitopes available only in a correctly folded protein structure, which would be lost in these experiments.

Patient immune reactions were also detected against the following proteins [cf. ref. 12]:

- Spot 10 – protein elongation factor EF-Tu (8/17)
- Spot 19 – ribosomal proteins S1 (5/17)
- Spot 12 – ribosomal protein L7/L12 (7/17)

Besides these known proteins, several new immunoreactive proteins were detected with frequencies ranging from 11/17 down to 1/17. The MW and pI characteristics of these proteins are shown in Table II. In addition, in a few cases, further analysis was performed by N-terminal amino acid sequencing supplemented with database homology searches.

Spot microsequencing

2D maps were prepared as described above, starting from 1mg total EB protein per run, followed by blotting onto polyvinylidene difluoride membranes (BioRad PVDF membranes 20 x 20 cm, 0.2 micron pore size), as in reference 20. The blots were stained with 0.1% (w/v)

Coomassie Brilliant Blue R250 in 50% aqueous methanol for 5 minutes, and de-stained in 40% methanol, 10% acetic acid. Membranes were dried at 37°C and stored at -20°C for further analysis. Selected protein spots were cut out and submitted to amino acid sequencing by Edman degradation using an automatic Protein/Peptide Sequencer (mod 470A; Applied Biosystem Inc.) connected on-line with a phenylthiohydantoin-amino acid analyser model 120A and a control/Data Module model 900A (Applied Biosystems Inc.). Typically 3 or 4 equivalent spots from similar blots were used, according to the estimated relative molar amount of protein in the spot.

The results of the sequencing are shown in Table III on page 16.

10 Computer analysis of sequences

Using the N-terminal sequence data, database searches for protein similarity were performed using the BLAST program [21] available from NCBI [<http://www.ncbi.nlm.nih.gov>] and programs of the GCG software (Wisconsin Package Version 9.0) [22]. Theoretical pI and MW values were calculated by the pI/MW computer program available from the ExPASy internet server [<http://www.expasy.ch>].

In addition to the usual databases, the genomic sequencing data of the *C.trachomatis* D/UW-3/Cx strain provided by the Chlamydia Genome Project [<http://chlamydia-www.berkeley.edu:4231>] was searched. Although the present study used a *C.trachomatis* serovar L2 strain (lymphogranuloma biovar), which has a different pathogenicity phenotype, several protein sequences could be safely correlated to the serovar D genes.

These searches with N-terminal data allowed the correlation of seven immunoreactive spots to known sequences (in addition to the seven noted above):

- Spot 15: predicted to be a periplasmic peptidase (currently annotated in the serovar D genomic database as *htrA*).
- Spots 18 & 46: predicted to be an outer membrane protein (currently annotated in the genomic database as *ompB*).
- Spot 21: Although the amino acid sequence does not match any previously-described proteins, it shows homology to an internal sequence from EF-Tu. This protein may be a breakdown or processing product of EF-Tu, or a variant.
- Spot 24: the RNA polymerase alpha subunit (*rpoA*)

- Spot 25: homologous to bacterial leucine peptidases (currently annotated in the genomic database as *pepA*).
- Spot 38: predicted to be a GTP-binding protein (currently annotated as *ychF*).

5 The N-terminal sequences of spots 26, 31 and 33 do not match any database sequences, including the published serovar D sequence.

Table IV shows a summary of identifications (some putative) of several immunoreactive antigens, which were obtained either by comparison with previous 2D mapping data, or by homology searches with the N-terminal sequencing data obtained above.

Proteins of particular interest

10 Of particular interest are the following proteins identified from the immunoblots:

Spot 24

This spot is believed to be the alpha chain of the *C. trachomatis* RNA polymerase (gi620029), based on its MW/pI position, and on its N-terminus sequence. Although the RNAP alpha chain has previously been described [23], it has never been reported as a chlamydial immunogen.

15 Four patients showed reactivity to this protein, demonstrating that it is immunogenic in humans as a consequence of chlamydial infection. Whilst the intracellular parasitic nature of *Chlamydia* means that it can generally evade antibody-mediated immune responses, the antibody reactivity demonstrated above indicates that the immune system does encounter these proteins during natural infection, and the formation of antibodies may, for instance, also help

20 to prime the T-cell-mediated immune responses.

Spots 18 & 46

Spots 18 and 46 appear to be homologous to the *ompB* gene in the serotype D genome, annotated as encoding a putative outer membrane protein. The N-terminal sequences and pI & MW values (at least for spot 18 – 5.08/34.09 vs predicted theoretical values of 5.06/34.5) are

25 in agreement with the expected properties of an *ompB* gene product, after cleavage of the predicted N-terminal signal peptide.

It has also been found that both spot 46 and 18 are present in a 2D electrophoretic map of a purified preparation of chlamydial outer membrane complex, which also supports the view that spots 18 and 46 represent the homologs of the serotype D *ompB* gene.

The reason why this protein appears as two distinct electrophoretic species was not investigated, but a spot shift of this type is usually associated to a variation of amino acid composition, either due to amino acid sequence variation, and/or to true or artefactual derivatisation of some amino acid residues.

- 5 Five patients showed reactivity to this protein, demonstrating that it is immunogenic in humans as a consequence of chlamydial infection.

Spot 25

- 10 This spot is believed to be an aminopeptidase, based on its MW/pI position and on its N-terminus sequence (both in comparison with the published serovar D sequence *pepA*). Four patients showed reactivity to this protein, demonstrating that it is immunogenic in humans as a consequence of chlamydial infection.

Spot 38

- 15 This spot is believed to be a GTP-binding protein, based on its MW/pI position and on its N-terminus sequence (both in comparison with the published serovar D sequence *ychF*). Two patients showed reactivity to this protein, demonstrating that it is immunogenic in humans as a consequence of chlamydial infection.

Spot 15

- 20 This spot is believed to be a stress-induced protease, based on its MW/pI position and on its N-terminus sequence (both in comparison with the published serovar D sequence *htrA*). Seven patients showed reactivity to this protein, demonstrating that it is immunogenic in humans as a consequence of chlamydial infection.

Spot 8

- 25 Nine patients showed reactivity towards protein spot 8, which could not be characterised by N-terminal sequencing. It does, however, have the following 'constellation type 2' amino acid composition (molar percentages):

aa	%	aa	%	aa	%	aa	%
Ala	6.5	Gly	22.5	Lys	3.7	Ser	13.7
Arg	3.5	His	0.5	Met	0.5	Thr	5.1
Asx	8.4	Ile	3.7	Phe	2.8	Tyr	2.2
Glx	12.5	Leu	6.7	Pro	3.4	Val	4.3

Cys and Trp are not determined in this type of analysis, and it is not possible to distinguish between Glu/Gln and Asp/Asn.

Inability to obtain N-terminal sequence, despite repeated attempts, suggests that the N-terminal residue is blocked due to some form of modification (*eg.* a lipoprotein).

- 5 Modification is often a characteristic of membrane-associated proteins in eukaryotes, but is also a characteristic of outer surface proteins or secreted proteins in bacterial species (*eg.* lipoproteins [24], mycoplasma outer membrane proteins [25], the FHA virulence factor of *B.pertussis* [26] *etc.*).

Spot 12

- 10 This spot is believed to be due to the ribosomal protein L7/L12. Seven patients showed reactivity to this protein, demonstrating that it is immunogenic in humans as a consequence of chlamydial infection. Although this protein has previously been described in chlamydia [accession number P38001] , it has never been reported as a chlamydial immunogen. It has, however, been described as an immunogen in Brucella infections [27,28].

15 Spot 19

This spot is believed to be due to the ribosomal protein S1. Five patients showed reactivity to this protein, demonstrating that it is immunogenic in humans as a consequence of chlamydial infection. Although this protein has previously been described in chlamydia [accession number P38016] , it has never been reported as an immunogen.

20 Spot 10

This spot is believed to be due to the protein synthesis elongation factor EF-Tu. Eight patients showed reactivity to this protein, demonstrating that it is immunogenic in humans as a consequence of chlamydial infection. Although the chlamydial EF-Tu has previously been described [accession number P26622] , it has never been reported as an immunogen.

25 Spots 10, 12, 15, 19 & 24

- Given the importance, in chronic infections, of a possible previous sensitisation to conserved microbial antigens that may trigger immunopathogenic reactions, it is noteworthy that several of these new immunoreactive antigens belong to conserved families of bacterial proteins: four (23%) sera reacted with spot 24 (the alpha subunit of the RNA polymerase); five (29%)
30 recognised spot 19 (ribosomal protein S1); eight (47%) recognised spot 10 (EF-Tu); seven (41%) recognised spot 15 (putative stress-induced protease of the HtrA (S2C peptidase)

family); and seven sera (41%) recognised spot 12 (the ribosomal protein L7/L12). In the group of sera used in this study, 12/17 (70.6%) reacted with at least one of these five antigens and, including the hsp60 and hsp70 antigens, all sera had antibodies reacting with between 1 and 7 (average 3.7) chlamydial proteins which have homologs in other bacteria.

- 5 Theories which postulate a role for immunological sensitisation mechanisms in chlamydial pathology, as described for the hsp60 GroEL-like antigen [29], should in fact be extended to several other common bacterial antigens, which may be immunogenic in other bacterial infections. For instance the protein elongation factor EF-Tu is immunogenic during the acute phase of infection with *Haemophilus influenzae*, and both L7/L12 and the HtrA stress-induced
- 10 protease homologues are immunogenic in *Brucella* infections. In the case of EF-Tu, the abundance of this protein in the bacterial cell may favour its "visibility" by the immune system. It should be noted, however, that EF-Tu has been described as associated to outer membrane and periplasmic cell fractions [30], and more recently data suggest that EF-Tu, in addition to its function in peptide elongation, has also a chaperone activity implicated in
- 15 protein folding and protection from stress [31]. Particularly intriguing is the response to the L7/L12 ribosomal protein, since in *Brucella melitensis* infections the homologous L7/L12 antigen induces a DTH cell-mediated response [27]. Furthermore vaccination of BALB/c mice with L7/L12 was shown to give protection against infection by *B.abortus* [32]. The unexpected finding that antibodies to L7/L12 are fairly frequent in patients infected by
- 20 *C.trachomatis* suggests that perhaps further attention should be paid to this antigens also in chlamydia-induced disease.

Spots 5, 6, 7, & 9

These proteins, whilst not yet correlated with any available genome sequence, and not yet having been sequenced, are of obvious interest given their prevalence (>50%) in the sera

25 tested.

It will be understood that the invention is described above by way of example only and modifications may be made whilst remaining within the scope and spirit of the invention.

TABLE I – SUMMARY OF PATIENT SERA

The letters in the first column correspond to those given in Table II. The codes in the second column refer to the original serum collection. The pathology associated with each patient is broadly indicated as cervicitis (lower genital tract infection), PID or sterility (secondary to infection). All sera were characterised by MIF assay with purified L2 elementary bodies. The MIF titre given in the table is the highest two-fold dilution which gave a positive signal. The 'Best Dilution' is the dilution which was found to give minimum background without loss of signal on weaker spots.

Serum ID in Table II	Original Serum ID	Pathology	MIF titre	Best Dilution
A	JO45/7931 BB	cervicitis	256	1:5000
B	JO28/7935 BB	cervicitis	16	1:1500
C	JO29/7936 BB	cervicitis	16	1:2000
D	JO51/7997 BB	cervicitis	256	1:5000
E	hs-C (Bologna)	P.I.D.	256	1:5000
F	14293 BB	P.I.D.	1024	1:10000
G	hs-B (Bologna)	P.I.D.	32	1:2000
H	JO6/7942BB	P.I.D.	256	1:5000
I	JO17/7953 BB	P.I.D.	256	1:5000
J	JO43/7989	P.I.D.	256	1:5000
K	JO20-/7956 BB	P.I.D.	256	1:5000
L	JO42/7988 BB	P.I.D.	256	1:5000
M	JO41/7987 BB	P.I.D.	256	1:5000
N	JO31/7977 BB	P.I.D.	256	1:5000
O	13839 BB	P.I.D.	256	1:5000
P	JO35/7934 BB	sterility	64	1:2500
Q	JO52/7933 BB	sterility	64	1:2500

TABLE II - PATIENT REACTIVITY WITH PROTEIN SPOTS

Spot #	pl	MW	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	FREQ
1	complex		+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	17
2	5.2-5.3	59.7	+	+	+	+	+	+	+	+	+	+	+			+	+	+	+	15
3	4.6-4.9	40	+	+	+	+	+	+	+		+	+		+		+	+		+	13
4	4.92-5.04	70.5	+	+	+	+	+	+			+	+	+		+				+	11
5	5.09	36.6	+	+	+	+	+	+			+	+				+		+	+	11
6	6.34	46.2-50	+			+	+	+			+	+	+			+		+	+	10
7	6.59	46.2-50.2	+			+	+	+			+	+	+			+		+	+	10
8	4.96	36.6	+		+		+	+			+	+				+		+	+	9
9	6.36	37.7-39.4	+	+			+	+			+	+	+					+	+	9
10	5.44-5.64	42.2	+	+	+		+		+				+					+	+	8
11	6.66	26.1		+			+	+			+		+			+	+		+	8
12	4.80	15.8		+		+	+				+		+	+						6
13	6.1	37.4-39.2	+	+			+						+			+		+	+	7
14	6.24	47.9	+				+	+			+	+	+					+		7
15	5.89	48.4	+	+	+		+	+					+					+		7
16	6.15	46-50	+				+				+	+	+					+	+	7
17	5.92	25.3		+			+	+								+	+		+	6
18	5.08	34.09			+	+		+					+						+	5
19	5.14-5.28	69	+		+	+		+					+							5
20	5.44	26.2		+			+									+	+		+	5
21	5.27	40.5		+	+	+							+						+	5
22	4.81	46.3	+		+								+	+		+				5
23	4.97	34.2			+			+					+							4
24	5.32	40.5			+	+							+						+	4
25	5.97	47.6	+	+														+	+	4
26	5.68	48.6	+					+					+						+	4
27	6.29-6.42	124.5	+				+					+						+		4
28	5.39	25.5					+	+								+				3
29	5.1	28.7					+				+					+				3
30	4.8	36.7			+		+												+	3
31	5.43	40.4				+							+						+	3
32	5.2-5.37	62.4			+			+					+							3
33	6.64	25.4					+				+									2
34	4.79	28.1				+													+	2
35	4.82	29.5				+													+	2
36	6.55	37.5					+					+								2
37	5.14	40.3		+															+	2
38	5.23	40.1																+	+	2
39	4.69	45.7		+										+						2
40	6.89	50										+						+		2
41	6.39-6.55	105	+				+													2
42	4.57	20.3									+									1
43	4.72	26.5					+													1
44	7.6	26.95											+							1
45	6.9	29.7														+				1
46	5.19	33.4				+														1
47	6.99	35.8											+							1
48	6.54	35.8											+							1
49	5.44	39.0																	+	1
50	5.37	41.0																	+	1
51	7.59	42.6											+							1
52	8.73	49.2					+									+				1
53	7.98	49.4			+															1
54	7.4	50.2											+							1
55	7.4	51.5											+							1

TABLE III – N-TERMINAL SEQUENCES OF PROTEINS

Spot #	N-terminal sequence
10	SKETFQRNK
12	TTESLETLVE
15	LAVSSGDQEVSQEDLLKE
18	XPAGNPAFPVIP
21	AKTRTLKGDG
24	SDSSHNLLYNK
25	VLLYSQASWDQRSKADAL
26	KAVYVQD(A/Q)E(V/D)Q
31	KDxxTNGQR
33	MSKGGQxD(Y/G)
38	XQXENGIVGL
46	MPAGNPAFPVIP

TABLE IV – IDENTIFICATION OF ANTIGENS

“CT-D gene” refers to the gene name from reference 2 and gives the names of genes likely to encode homologue proteins in *C. trachomatis* D. Theoretical pI/MW values in the last column, to be compared to the experimental values, were calculated from CT-D gene sequences.

spot	Map location	N-terminal AA seq	Annotation	CT-D gene	Predicted pI/MW
1	OMP2 cluster	-	OMP2	<i>omcB</i>	7.65-7.92/54.5-58.7
2	5.2-5.3/59.7	VA(D/K)NI(K/F)YNEE	GroEL-like	<i>groEL1</i>	5.11/58.1
3	4.6-4.9/40	LPVGN	MOMP	<i>ompA</i>	4.69/40.3
4	4.92-5.04/70.5	SEKRK(S/A)N(K/S)....	DnaK-like	<i>dnaK</i>	4.88/70.7
10	5.44-5.64/42.2	SKETFQRNK	EF-Tu	<i>tufA</i>	5.36/43.1
12	4.80/15.8	TTESLETLVE	Ribosomal protein L7/12	<i>r17</i>	5.09/13.5
15	5.89/48.4	LAVSSGDQEVSQEDLLKE	stress induced protease	<i>htrA</i>	5.83/49.5
18	5.08/34.09	XPAGNPAFPVIP	outer membrane protein	<i>ompB</i>	5.06/34.5
19	5.14-5.28/69	Not determined	Ribosomal protein S1	<i>rs1</i>	5.17/63.6
21	5.27/40.5	AKTRTLKGDG	EF-Tu related peptide?	-	-
24	5.32/40.5	SDSSHNLLYNK	RNAP alpha chain	<i>rpoA</i>	5.34/41.7
25	5.97/47.6	VLLYSQASWDQRSKADAL	Aminopeptidase	<i>pepA</i>	5.74/54.0
26	5.68/48.6	KAVYVQD(A/Q)E(V/D)Q	Not identified	-	-
31	5.43/40.4	KDxxTNGQR	Not identified	-	-
33	6.64/25.4	MSKGGQxD(Y/G)	Not identified	-	-
38	5.23/40.1	XQXENGIVGL	GTP-binding protein	<i>ychF</i>	5.16/39.5
46	5.19/33.4	MPAGNPAFPVIP	outer membrane protein	<i>ompB</i>	5.06/34.5

REFERENCES (the contents of which are incorporated herein in their entirety)

- 1 Raulston (1995) Chlamydial envelope components and pathogen-host cell interactions. *Mol Microbiol* 15(4):607-616.
- 2 Stephens *et al.* (1998) Genome Sequence of an Obligate Intracellular Pathogen of Humans: *Chlamydia trachomatis*. *Science* 282:754-759.
- 3 Ward (1995) The immunobiology and immunopathology of chlamydial infections. *Apmis*. 103:769-96.
- 4 Moulder (1991) Interaction of *Chlamydiae* and host cells *in vitro*. *Microbiol Rev* 55(1):143-190.
- 5 Comanducci *et al.* (1994) Humoral immune response to plasmid protein pgp3 in patients with *Chlamydia trachomatis* infection. *Infect Immun* 62(12):5491-5497.
- 6 EP-A-0499681
- 7 WO95/28487
- 8 Murdin *et al.* (1993) *Infect Immun* 61:4406-4414
- 9 Cerrone *et al.* (1991) Cloning and sequence of the gene for heat shock protein 60 from *Chlamydia trachomatis* and immunological reactivity of the protein. *Infect Immun* 59(1):79-90.
- 10 Raulston *et al.* (1993) Molecular characterization and outer membrane association of a *Chlamydia trachomatis* protein related to the hsp70 family of proteins. *J. Biol. Chem.* 268:23139-23147.
- 11 Wang & Grayston (1970) Immunologic relationship between genital TRIC, lymphogranuloma venereum, and related organisms in a new microtiter indirect immunofluorescence test. *Am. J. Ophthalmol.* 70:367-374
- 12 Bini *et al.* (1996) Mapping of *Chlamydia trachomatis* proteins by immobililine-polyacrylamide two-dimensional electrophoresis: spot identification by N-terminal sequencing and immunoblotting. *Electrophoresis* 17:185-190.
- 13 Schacter & Wyrick (1994) Culture and isolation of *Chlamydia trachomatis*. *Meth Enzymol* 236:377-390
- 14 Görg *et al.* (1988) The current state of two-dimensional electrophoresis with immobilized pH gradients. *Electrophoresis* 9:531-546
- 15 Hochstrasser *et al.* (1988) Methods for increasing the resolution of two-dimensional protein electrophoresis. *Anal. Biochem.* 173:424-435.
- 16 Oakley *et al.* (1980) A simplified ultrasensitive silver stain for detecting proteins in polyacrylamide gels. *Anal. Biochem.* 105:361-363.
- 17 Bjellqvist *et al.* (1993) A nonlinear wide-range immobilized pH gradient for two-dimensional electrophoresis and its definition in a relevant pH scale. *Electrophoresis* 14: 1357-1365

- 18 Towbin *et al.* (1979) Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci USA* 76: 4350-4354.
- 19 Magi *et al.* Chapter entitled "Immunoaffinity identification of 2-DE separated proteins" in *Methods in Molecular Biology*, 2-D Proteome analysis protocols. AJ Link (*ed.*), Humana Press, Totowa, NJ, USA.
- 20 Matsudaira (1987) Sequence from picomole quantities of proteins electroblotted onto polyvinylidene difluoride membranes. *J Biol Chem* 262:10035-10038.
- 21 Altschul *et al.* (1990) Basic local alignment search tool. *J Mol Biol* 215:403-410.
- 22 Devereux *et al.* (1984) A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res* 12: 387-395.
- 23 Gu *et al.* (1995) *Chlamydia trachomatis* RNA polymerase alpha subunit: sequence and structural analysis. *J Bacteriol* 177:2594-2601.
- 24 Nyström *et al.* (1992) Membrane protein acylation. Preference for exogenous myristic acid or endogenous saturated chains in *Acholeplasma laidlawii*. *Eur J Biochem* 204:231-40
- 25 Jan *et al.* (1995) Acylation and immunological properties of *Mycoplasma gallisepticum* membrane proteins. *Res Microbiol* 146:739-50
- 26 Lambert-Buisine *et al.* (1998) N-terminal characterization of the Bordetella pertussis filamentous haemagglutinin. *Mol Microbiol* 28:1283-93
- 27 Bachrach *et al.* (1994) Brucella ribosomal protein L7/L12 is a major component in the antigenicity of brucellin INRA for delayed-type hypersensitivity in Brucella-sensitized guinea pigs. *Infect. Immun.* 62:5361-5366
- 28 Roop *et al.* (1994) Identification of an immunoreactive *Brucella abortus* HtrA stress response protein homolog. *Infect. Immun.* 62:1000-1007
- 29 Morrison *et al.* (1989) Chlamydial disease pathogenesis: the 57-kD chlamydial hypersensitivity antigen is a stress response protein *J.Exp.Med.* 170:1271-1283. Similarly: Morrison *et al.* (1989) Chlamydial disease pathogenesis: ocular hypersensitivity elicited by a genus-specific 57-kD protein. *J.Exp.Med.* 169:663-675
- 30 Marques *et al.* (1998) Mapping and Identification of the Major Cell Wall-Associated Components of *Mycobacterium leprae*. *Infect. Immun.* 66: 2625-2631.
- 31 Caldas *et al.* (1998) Chaperone properties of bacterial elongation factor EF-Tu. *J Biol. Chem.* 273:11478-82
- 32 Oliveira & Splitter (1996) Immunization of mice with recombinant L7/L12 ribosomal protein confers protection against *Brucella abortus* infection. *Vaccine* 14:959-62

CLAIMS

1. A *C.trachomatis* protein having the MW and pI characteristics of protein 5, 6, 7, 8, 9, 11, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, or 55, as set out in Table II
5 on page 15.
2. A protein according to claim 1 having, in the L2 strain of *C.trachomatis*, an N-terminal amino acid sequence disclosed in Table III on page 16.
3. A protein having 50% or more sequence identity to a protein according to claim 1
4. A protein comprising a fragment of at least 7 consecutive amino acids of a *C.trachomatis*
10 protein according to claim 1
5. An antibody which binds to a protein according to any one of claims 1 to 4.
6. Nucleic acid encoding a protein according to any one of claims 1 to 4.
7. Nucleic acid having 50% or more sequence identity to the nucleic acid of claim 6.
8. Nucleic acid which can hybridise to the nucleic acid of claim 6.
- 15 9. Nucleic acid comprising a fragment of 10 or more consecutive nucleotides of the nucleic acid according to claim 6.
10. A vector comprising nucleic acid according to claim 6.
11. A host cell transformed with a vector according to claim 10.
12. A composition comprising a protein according to any one of claims 1 to 4, an antibody
20 according to claim 5, and/or nucleic acid according to any one of claims 6 to 9.
13. A protein according to any one of claims 1 to 4, an antibody according to claim 5, and/or nucleic acid according to any one of claims 6 to 9 for use as a medicament or as a diagnostic reagent.
14. A *C.trachomatis* protein having the MW and pI characteristics of protein 5, 6, 7, 8, 9, 10,
25 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, or 55 (as set out in Table II on page 15) for use as a chlamydial immunogen.

15. The use of a protein according to any one of claims 1 to 4, an antibody according to claim 5, and/or nucleic acid according to any one of claims 6 to 9, in the manufacture of a medicament for treating or preventing infection due to *Chlamydia*.
- 5 16. The use of a protein according to any one of claims 1 to 4, an antibody according to claim 5, and/or nucleic acid according to any one of claims 6 to 9, in the manufacture of a diagnostic reagent for detecting the presence of *Chlamydia* or of antibodies raised against *Chlamydia*.
- 10 17. The use of a protein according to any one of claims 1 to 4, an antibody according to claim 5, and/or nucleic acid according to any one of claims 6 to 9, in the manufacture of a reagent which can raise antibodies against *Chlamydia*.
18. A method of treating a patient, comprising administering to the patient a therapeutically effective amount of a protein according to any one of claims 1 to 4, an antibody according to claim 5, and/or nucleic acid according to any one of claims 6 to 9.
- 15 19. A process for producing a protein according to any one of claims 1 to 4, comprising the step of culturing a host cell according to claim 11 under conditions which induce protein expression.
- 20 20. A process for producing a protein according to any one of claims 1 to 4 or nucleic acid according to any one of claims 6 to 9, wherein the protein or nucleic acid is synthesised in part or in whole using chemical means.
- 20 21. A process for detecting nucleic acid according to claim any one of claims 6 to 9, comprising the steps of: (a) contacting a nucleic acid probe with a biological sample under hybridising conditions to form duplexes; and (b) detecting said duplexes.
- 25 22. A process for detecting a protein according to any one of claims 1 to 4, comprising the steps of: (a) contacting an antibody according to claim 5 with a biological sample under conditions suitable for the formation of an antibody-antigen complexes; and (b) detecting said complexes.
- 30 23. A process for detecting an antibody according to claim 5, comprising the steps of: (a) contacting a protein according to any one of claims 1 to 4 with a biological sample under conditions suitable for the formation of an antibody-antigen complexes; and (b) detecting said complexes.
24. A kit comprising reagents suitable for use in a process according to any one of claims 21 to 23.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/295, 16/12, C12N 15/31, 5/10, A61K 38/16, 39/118, 39/395, 48/00, G01N 33/569, C12P 21/00, C12Q 1/68		A3	(11) International Publication Number: WO 00/37494 (43) International Publication Date: 29 June 2000 (29.06.00)
(21) International Application Number: PCT/IB99/02065 (22) International Filing Date: 17 December 1999 (17.12.99) (30) Priority Data: 9828000.1 18 December 1998 (18.12.98) GB (71) Applicant (for all designated States except US): CHIRON S.P.A. [IT/IT]; Via Fiorentina, 1, I-53100 Siena (IT). (72) Inventor; and (75) Inventor/Applicant (for US only): RATTI, Giulio [IT/IT]; Iris Research Centre, Via Fiorentina, 1, I-53100 Siena (IT). (74) Agent: HALLYBONE, Huw, George; Carpmaels & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report. (88) Date of publication of the international search report: 12 October 2000 (12.10.00)	

(54) Title: CHLAMYDIA TRACHOMATIS ANTIGENS


(57) Abstract

Proteins encoded by *Chlamydia trachoma-
tis* which are immunogenic in humans as a con-
sequence of infection have been identified using
Western blots of two-dimensional electrophoretic
maps. Several known immunogens were identified,
as were proteins not previously known to be im-
munogens, and proteins not previously reported as
expressed gene products.



1/3

FIG. 1



2/3

FIG. 2(A)





FIG. 2(B)



3/3

FIG. 2(C)





FIG. 2(D)



I hereby certify that this paper is being deposited with the United States Postal Service as first class mail in an envelope addressed: BOX MISSING PARTS, Assistant Commissioner of Patents and Trademarks, Washington, D.C. 20231 on February 22, 2002.

Gyne Riser
Gyne Riser

02.22.02
Date

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: GIULIO RATTI
International Filing No.: PCT/IB99/02065 Examiner: Unassigned
U.S Serial No.: 09/868,293 Group Art Unit: Unassigned
International Filing Date: December 17, 1999
For: CHLAMYDIA TRACHOMATIS ANTIGENS

POWER OF ATTORNEY

Assistant Commissioner of Patents
and Trademarks
Washington, D.C. 20231

Sir:

Pursuant to 37 C.F.R. §1.36, Chiron Corporation, the Assignee of the above-identified patent application hereby appoints the following attorneys and agents to prosecute this application and to transact all business in the Patent and Trademark Office in connection therewith.

Lisa A. Alexander - Reg. No. 41,576
Robert P. Blackburn - Reg. No. 30,447
Anne S. Dollard - Reg. No. 43,935
Joseph H. Guth - Reg. No. 31,261
Rebecca M. Hale - Reg. No. 45,680
Alisa A. Harbin - Reg. No. 33,895
Charlene A. Launer - Reg. No. 33,035

David P. Lentini - Reg. No. 33,944
Kimberlin L. Morley - Reg. No. 35,391

Please direct all correspondence and telephone calls regarding this application to the following:

Chiron Corporation
Intellectual Property - R440
P.O. Box 8097
Emeryville, CA 94662-8097
(510) 923-3179

CHIRON CORPORATION

Dated: *Feb 22, 2002*

By: *Rebecca M. Hale*
Rebecca M. Hale
Assistant Secretary

DECLARATION FOR PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: CHLAMYDIA TRACHOMATIS ANTIGENS

the specification of which (check one) ☐ is attached hereto ☒ was filed internationally on December 17, 1999, as Application Serial No. PCT/IB99/02065, and was amended on (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s)		Day/Month/Year Filed	Priority Claimed	
Number	Country		Yes	No
GB 9828000.1		December 18, 1998	X	

I hereby claim the benefit under Title 35, United States Code, §120 and/or §119(e) of any United States application(s) and/or provisional applications listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, §1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

Application Serial No.	Filing Date	Status
		Patented, Pending, Abandoned

I hereby declare that all statements herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements and the like so made are punishable by fine or imprisonment or both, under Section 1001 of Title 18 of the United States code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1641-102 sequence listing.TXT

SEQUENCE LISTING

```

<110>          CHIRON SpA
<120>          CHLAMYDIA TRACHOMATIS ANTIGENS
<140>          PCT/IB99/02065
<141>          1999-12-17
<150>          GB-9828000.1
<151>          1998-12-18
<160>          15
<170>          SeqWin99, version 1.02
<210>          1
<211>          9
<212>          PRT
<213>          Chlamydia trachomatis
<400>          1
Ser Lys Glu Thr Phe Gln Arg Asn Lys
1          5
<210>          2
<211>          10
<212>          PRT
<213>          Chlamydia trachomatis
<400>          2
Thr Thr Glu Ser Leu Glu Thr Leu Val Glu
1          5          10
<210>          3
<211>          18
<212>          PRT
<213>          Chlamydia trachomatis
<400>          3
Leu Ala Val Ser Ser Gly Asp Gln Glu Val Ser Gln Glu Asp Leu Leu
1          5          10          15
Lys Glu
<210>          4
<211>          12
<212>          PRT
<213>          Chlamydia trachomatis
<220>          Xaa
<221>          1
<222>          1
<223>          Xaa is any amino acid
<400>          4
Xaa Pro Ala Gly Asn Pro Ala Phe Pro Val Ile Pro
1          5          10
<210>          5
<211>          10
<212>          PRT

```

SECRET

<400> 5
Ala Lys Thr Arg Thr Leu Lys Gly Asp Gly
1 5 10

<400> 6
Ser Asp Ser Ser His Asn Leu Leu Tyr Asn Lys
1 5 10

<400> 7
Val Leu Leu Tyr Ser Gln Ala Ser Trp Asp Gln Arg Ser Lys Ala Asp
1 5 10 15

<210>	8
<211>	11
<212>	PRT
<213>	Chlamydia trachomatis

```
<220>
<221> xaa
<222> 10
<223> xaa is val or Asp
```

<210>	9
<211>	9
<212>	PRT
<213>	Chlamydia trachomatis

```
<220>
<221> xaa
<222> 4
<223> xaa is any amino acid
```

Page 2

1641-102 sequence listing.TXT

1

5

```
<210> 10
<211> 10
<212> PRT
<213> Chlamydia trachomatis
```

```
<220>
<221>    Xaa
<222>    8
<223>    Xaa is any amino acid
```

```
<220>
<221>    xaa
<222>    10
<223>    xaa is Tyr or Gly
```

<400> 10
Met Ser Lys Gly Gly Gln Thr Xaa Asp Xaa
1 5 10

```
<210> 11
<211> 10
<212> PRT
<213> Chlamydia trachomatis
```

```
<220>
<221>    xaa
<222>    1
<223>    xaa is any amino acid
```

```
<220>
<221>    Xaa
<222>    3
<223>    Xaa is any amino acid
```

<400> 11
xaa Gln xaa Glu Asn Gly Ile val Gly Leu
1 5 10

<210> 12
<211> 12
<212> PRT
<213> Chlamydia trachomatis

<400> 12
Met Pro Ala Gly Asn Pro Ala Phe Pro Val Ile Pro
1 5 10

```
<210> 13
<211> 10
<212> PRT
<213> Chlamydia trachomatis
```

```
<220>
<221>    xaa
<222>    3
<223>    xaa is Asp or Lys
```

```
<220>
<221>    xaa
<222>    6
<223>    xaa is Lys or Phe
```

1641-102 Sequence listing.TXT

<400> 13
Val Ala Xaa Asn Ile Xaa Tyr Asn Glu Glu
1 5 10

<210> 14
<211> 5
<212> PRT
<213> Chlamydia trachomatis

<400> 14
Leu Pro Val Gly Asn
1 5

<210> 15
<211> 8
<212> PRT
<213> Chlamydia trachomatis

<220>
<221> Xaa
<222> 6
<223> Xaa is Ser or Ala

<220>
<221> Xaa
<222> 8
<223> Xaa is Lys or Ser

<400> 15
Ser Glu Lys Arg Lys Xaa Asn Xaa
1 5